

General Immunization Information

	Schedule	Vaccine	Day/mo/yr. dose was given
1.	Diphtheria, Pertussis and Tetanus (DPT) or Diphtheria/Tetanus (DT) schedule DPT <ul style="list-style-type: none"> • Given to children 6 wks to 6 yrs of age • 3 doses 2 mos. Apart • 4th dose 1 yr. After the 3rd. • Booster doses every 10 yrs. DT <ul style="list-style-type: none"> • 2 doses 1 mo. apart • 3rd dose 1 yr. After 2nd dose • Booster doses every 10 yrs. 	Diphtheria/ Pertussis/ Tetanus (DPT) or (DT)	1 st 2 nd 3 rd 4 th 5 th 6 th
2.	Polio Schedule <ul style="list-style-type: none"> • Given to children 6 wks. to 19 yrs. of age • 2 doses 6-8 wks. apart • 3rd dose 6-12 months after the 2nd. • 4th dose given upon entering school • Booster dose given in some countries 	Polio	1 st 2 nd 3 rd 4 th 5 th
3.	Measles (10-day or Red Measles), Mumps and/or Rubella (3 day German Measles Schedule): <ul style="list-style-type: none"> • Given to children 1 yr. of age and older • Booster given in some countries at 4-6 yrs. or 12 yrs. 	Measles Mumps Rubella	1 st 2 nd 1 st 2 nd 1 st 2 nd
4.	BCG (for TB) <ul style="list-style-type: none"> • Usually given at birth • Tuberculin test recommended for adults • One dose only 	BCG	1 st
5.	Hepatitis B (HBV) <ul style="list-style-type: none"> • Given at birth • 1 mo. • 6-12 mos. 	HBV	1 st 2 nd
6.	Hemophilus Influenza B (HIB) <ul style="list-style-type: none"> • Usually given at 2 mos., 4 mos., 6 mos., and 12-15 mos. 	HIB	1 st 2 nd 3 rd 4 th
7.	Hepatitis A (HAV) <ul style="list-style-type: none"> • Highly recommended after 5 yrs. • 3 doses – 2nd dose 1 mo. after 1st. • 3rd dose 6-12 mos. After 2nd. 	HAV	1 st 2 nd 3 rd

Permission is hereby given for emergency measures to be initiated in case of accident or sudden illness with the understanding that I will be notified. I certify that all information on this record is complete and correct.

Parent's signature.....Date.....